

The enclosed prescription medication is made by ImprimisRx, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. Imprimis Pharmaceuticals dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. Imprimis Pharmaceuticals does not compound copies of commercially available products. References available upon request.

ImprimisRx is trademarks of Harrow Health, Inc. ©2019 ImprimisRx. All rights reserved.
IMPO0375 03/19 As of 03.28.19

**Questions?
Contact us at:
(844) 446-6979**

Imprimis Rx
12264 El Camino Real, Suite 350
San Diego, CA 92130

imprimis Rx[®]

Product Name:

Prednisolone Acetate, Moxifloxacin
and Nepafenac

Ophthalmic Suspension

		Prednisolone Acetate	Moxifloxacin	Nepafenac
	Possible Indications	Postoperative Ophthalmic Inflammation Postoperative Eye Pain	Bacterial Conjunctivitis	Postoperative Ophthalmic Inflammation Postoperative Ophthalmic Pain
	Possible Adverse Effects	Accommodation Disturbance Conjunctival Hyperemia Decreased Visual Acuity Eye Infection Glaucoma Increased Intraocular Pressure Keratitis Mydriasis Optic Nerve Damage Subscapular Posterior Cataract Visual Field Defect Wound Healing Impairment	Conjunctival Irritation Decreased Visual Acuity Dysgeusia Eye Discharge Eye Irritation Eye Pain Eye Redness Eyelid Edema Headache Increased Lacrimation Keratitis Papillary Conjunctivitis Pyrexia Subconjunctival Hemorrhage Tearing Hypersensitivity Reaction	Abnormal Sensation in Eyes Increased Intraocular Pressure Anterior Chamber Eye Hemorrhage Burning Sensation Conjunctival Hyperemia Corneal Erosion Corneal Perforation Corneal Thinning Corneal Ulcer Epithelial Keratopathy Eye Irritation Eye Pain Eye Pruritus Eye Redness Headache Hypersensitivity Reaction Iritis Keratitis Prolonged Bleeding Stinging Sensation
	Potential Contraindications/Precautions	Infants Lactation Pregnancy Cataract Eye Infection Increased Intraocular Pressure	Infants Lactation Pregnancy	Lactation Pediatric Patients Pregnancy Corneal Disease Thrombocytopenia (bleeding disorders)
	Storage	Store at Room Temperature	20°C to 25°C (68°F to 77°F)	
	Auxiliary Labeling	Shake Well Before Use!	Store upright in original bottle	Protect from light.