

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs.

This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions.

ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

ImprimisRx is a registered trademark of Harrow Health, Inc. ©2020 ImprimisRx. All rights reserved. IMPO0376 Rev2 06/20  
As of 06.15.20

**Questions?  
Contact us at:  
(844) 446-6979**

ImprimisRx®  
12264 El Camino Real, Suite 350  
San Diego, CA 92130

**imprimis** Rx®

**Product Name:**

Prednisolone Acetate  
Nepafenac

---

Ophthalmic Suspension for Topical Use

**Prednisolone Acetate****Nepafenac****Possible Indications**

Postoperative Ophthalmic Inflammation  
Postoperative Eye Pain

Postoperative Ophthalmic Inflammation  
Postoperative Ophthalmic Pain

**Possible Adverse Effects**

Accommodation Disturbance  
Anterior Uveitis  
Blepharitis  
Conjunctival Hyperemia  
Conjunctivitis  
Corneal Perforation or Ulcer  
Decreased Visual Acuity  
Eye Infection  
Glaucoma  
Increased Intraocular Pressure  
Keratitis  
Mydriasis  
Optic Nerve Damage  
Visual Field Defect  
Wound Healing Impairment

Abnormal Sensation in Eyes  
Increased Intraocular Pressure  
Anterior Chamber Eye Hemorrhage  
Burning Sensation  
Conjunctival Hyperemia  
Corneal Erosion  
Corneal Perforation  
Corneal Thinning  
Corneal Ulcer  
Epithelial Keratopathy  
Eye Irritation  
Eye Pain  
Eye Pruritus  
Eye Redness  
Headache  
Hypersensitivity Reaction  
Iritis  
Keratitis  
Prolonged Bleeding  
Stinging Sensation

**Potential Contraindications/  
Precautions**

Infants  
Lactation  
Pregnancy  
Cataract  
Eye Infection  
Increased Intraocular Pressure

Lactation  
Pediatric Patients  
Pregnancy  
Corneal Disease  
Thrombocytopenia (bleeding disorders)

**Storage**

Store at Room Temperature

20°C to 25°C (68°F to 77°F)

**Auxiliary Labeling**

Shake Well Before Use!