

The enclosed prescription medication is made by ImprimisRx, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

ImprimisRx is trademarks of Harrow Health, Inc. ©2019 ImprimisRx. All rights reserved. IMPO0453 11/19

**Questions?
Contact us at:
(844) 446-6979**

imprimis Rx[®]

Product Name:

**Brimonidine
Dorzolamide**

Ophthalmic Solution
Preservative-Free

ImprimisRx
12264 El Camino Real, Suite 350
San Diego, CA 92130

		Brimonidine Tartrate	Dorzolamide	
	Possible Indications	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma	
	Possible Adverse Effects	<p>Oral dryness</p> <p>Ocular hyperemia,</p> <p>Burning and Stinging on Instillation</p> <p>Headache</p> <p>Blurred vision</p> <p>Foreign body sensation</p> <p>Fatigue/drowsiness</p> <p>Conjunctival follicles</p> <p>Ocular allergic reactions</p> <p>Ocular pruritis</p> <p>Corneal staining/erosion</p> <p>Photophobia</p> <p>Eyelid erythema</p> <p>Ocular ache/pain</p> <p>Upper respiratory symptoms</p> <p>Dizziness</p> <p>Blepharitis, ocular irritation</p> <p>Lid crusting</p> <p>Conjunctival hemorrhaging</p> <p>Abnormal taste, insomnia, depression, hypertension, anxiety, palpitations/arrhythmias, nasal dryness and syncope</p>	<p>Ocular burning, stinging, or discomfort immediately following ocular administration</p> <p>Bitter taste</p> <p>Superficial punctate keratitis</p> <p>Ocular allergic reaction</p> <p>Conjunctivitis and lid reactions</p> <p>Blurred vision</p> <p>Eye redness</p> <p>Tearing</p> <p>Dryness</p> <p>Photophobia</p> <p>Headache</p> <p>Nausea</p> <p>Asthenia/fatigue</p> <p>Skin rashes</p> <p>Urolithiasis</p> <p>Iridocyclitis</p>	
	Potential Contraindications/Precautions	<p>Infants</p> <p>Lactation</p> <p>Pregnancy</p> <p>Concomitant MAO Inhibitor Therapy</p>	<p>Neonates and Patients Less than 12 Years</p> <p>Lactation</p> <p>Pregnancy</p>	
	Storage	Store at room temperature	20°C to 25°C (68°F to 77°F)	
	Auxiliary Labeling	For professional use only.	Store upright in original bottle.	Protect from light.