

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

ImprimisRx is a registered trademark of Harrow IP, LLC. ©2022 ImprimisRx. All Rights Reserved. IMPO0637 Rev1 09/22



Product Name:

Tobramycin – Vancomycin

**Questions?
Contact us at:
(844) 446-6979**

Fortified Ophthalmic Solution

ImprimisRx®
1000 Aviara Parkway, Suite 220
Carlsbad, CA 92011

		Tobramycin	Vancomycin	
	Possible Indications	Treatment of infections of the eye caused by susceptible bacteria	Treatment of infections of the eye caused by susceptible bacteria	
	Possible Adverse Effects	<p>Ocular burning/stinging on instillation Lid itching and swelling Conjunctival erythema Anaphylactic reaction Stevens-Johnson syndrome Erythema multiforme</p> <p>The following are adverse reactions reported with systemic aminoglycosides:</p> <ul style="list-style-type: none"> • Neurotoxicity • Ototoxicity • Nephrotoxicity • Aminoglycosides may aggravate muscle weakness in patients with known or suspected neuromuscular disorders, such as myasthenia gravis or Parkinson's disease. 	<p>Ocular burning/stinging on instillation Lid itching and swelling Conjunctival erythema Anaphylactic reaction Stevens-Johnson syndrome</p> <p>The following are adverse reactions reported with systemic vancomycin administration:</p> <ul style="list-style-type: none"> • Nephrotoxicity • Onset of pseudomembranous colitis symptoms • Ototoxicity • Neutropenia <p>Hemorrhagic occlusive retinal vasculitis (HORV) may be associated with the use of intraocular vancomycin (including intravitreal, intracameral bolus, and irrigating solution containing vancomycin). Although the precise etiology of HORV remains unproven, each surgeon should weigh the potential risk of vancomycin against the risk of endophthalmitis.</p>	
	Potential Contraindications/Precautions	<p>Patients Less than 12 Years</p> <p>Nursing Mothers</p> <p>Pregnancy</p>	<p>Patients Less than 12 Years</p> <p>Nursing Mothers</p> <p>Pregnancy</p>	
	Storage	<p>Store under refrigeration</p> <p>2°C to 8°C (36°F to 46°F)</p>	<p>Store under refrigeration</p> <p>2°C to 8°C (36°F to 46°F)</p>	
	Auxiliary Labeling	Protect from light	Protect from light	