

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

ImprimisRx is a registered trademark of Harrow Health, Inc. ©2021 ImprimisRx. All rights reserved. INT0017Rev2 06/21

**Questions?
Contact us at:
(844) 446-6979**

imprimis Rx®

Product Name:

Ascorbic Acid
(Corn Protein Free)

Injection Solution
Preservative-Free

ImprimisRx®
12264 El Camino Real, Suite 350
San Diego, CA 92130

Ascorbic Acid

Possible Indications

Prevention and treatment of scurvy
Treatment of symptoms of mild deficiency

Possible Adverse Effects

Hyperoxaluria (with large doses)
Diarrhea
Dizziness
Fatigue
Flank pain
Flushing
Headache
Heartburn
Nausea
Vomiting

Potential Contraindications/ Precautions

Oxalate nephropathy/nephrolithiasis; elderly patients and pediatric patients <2 years of age may be at increased risk. Patients with diabetes mellitus should not take excessive doses for extended periods of time. Hemolysis has been reported in patients with glucose-6-phosphatase dehydrogenase (G6PD) deficiency and the risk for severe hemolysis may be increased during ascorbic acid therapy. Use with caution in patients with hemochromatosis. Use with caution in patients with renal impairment or patients prone to recurrent renal calculi; may have increased risk of developing acute or chronic oxalate nephropathy. Elderly and Pediatric: Use with caution in the elderly and children <2; may be at increased risk for oxalate nephropathy.

Storage

Store at Room Temperature 20°C to 25°C (68°F to 77°F).

Precautions

Bulk Package - Must dilute before use.
Pressure may build up inside vial.

Auxiliary Labeling

Keep all medicines out of the reach of children. Protect from light. Single-Use vial.
Discard unused portion. Throw away any medicine after the beyond use date.