The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

ImprimisRx is a registered trademark of Harrow Health, Inc. ©2020 ImprimisRx. All rights reserved. INT0033 06/20 As of 06.10.20

**Questions?**
**Contact us at:**
**(844) 446-6979**

ImprimisRx
12264 El Camino Real, Suite 350
San Diego, CA 92130

imprimis Rx®

**Product Name:**

Taurine

Injection Solution

## Taurine

| | |
|---|---|
| **Possible Indications** | Taurine is indicated in those with congestive heart failure (CHF), liver disease (hepatitis), retinal disorder, chemotherapy side effects, cirrhosis, cystic fibrosis, diabetes, fatigue, high blood pressure, mental performance, insomnia, psychosis and anemia. |
| **Possible Adverse Effects** | No noted side effects when dosed within normal limits. |
| **Potential Contraindications/ Precautions** | Recommended dosage if taurine is less than 3,000 mg/day. Dosage is different for each treatment. Several mediations may cause interactions with Taurine. Lithium may have a diuretic effect, Antiplatelets/ Anticoagulants may lower the clotting effect leading to bleeding, NSAIDs / Salicylates may lead to changed clotting times and bleeding. |
| **Storage** | Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) (See USP Controlled Room Temperature). |
| **Auxiliary Labeling** | |