

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound copies of commercially available products. References available upon request.

ImprimisRx is a registered trademark of Imprimis Pharmaceuticals, Inc.  
©2018 Imprimis Pharmaceuticals, Inc. All Rights Reserved. IMPO0306 9/18

**Questions?**  
**Contact us at:**  
**(844) 446-6979**

Imprimis Pharmaceuticals, Inc.  
12264 El Camino Real, Suite 350  
San Diego, CA 92130

**imprimis** Rx®

**Product Name:**

Dexamethasone Sodium Phosphate

Moxifloxacin HCl

Ketorolac

---

Intraocular Solution for Injection

	<b>Dexamethasone Sodium Phosphate</b>	<b>Moxifloxacin HCl</b>	<b>Ketorolac Tromethamine</b>	
<b>Possible Indications</b>	<ul style="list-style-type: none"> <li>Postoperative Inflammation</li> <li>Allergic Conjunctivitis</li> <li>Inflammation of Anterior Segment of Eye</li> <li>Keratitis, Optic Neuritis, Iritis</li> <li>Sympathetic Ophthalmia</li> </ul>	<ul style="list-style-type: none"> <li>Bacterial Conjunctivitis</li> </ul>	<ul style="list-style-type: none"> <li>Postoperative Eye Pain</li> <li>Transient Burning and Stinging in the Eyes</li> </ul>	
<b>Possible Adverse Effects</b>	<ul style="list-style-type: none"> <li>Conjunctival Irritation</li> <li>Burning and Stinging Sensation</li> <li>Fever</li> <li>Eye Irritation and Infection</li> <li>Airway Obstruction</li> <li>Anaphylaxis</li> <li>Angioedema</li> <li>Circulatory Shock</li> <li>Blindness</li> <li>Post-Injection Flare</li> <li>Wound Healing Suppression</li> <li>Anaphylaxis</li> <li>Corneal Perforation</li> <li>Headache</li> <li>Subcapsular Posterior Cataract</li> <li>Ocular Hypertension</li> <li>Optic Nerve Damage</li> <li>Visual Field Defect</li> </ul>	<ul style="list-style-type: none"> <li>Conjunctival Irritation</li> <li>Fever</li> <li>Eye Irritation</li> <li>Airway Obstruction</li> <li>Anaphylaxis</li> <li>Angioedema</li> <li>Circulatory Shock</li> <li>Dyspnea</li> <li>Facial Edema</li> <li>Fungal Superinfection</li> <li>Laryngeal Edema</li> <li>Loss of Consciousness</li> <li>Pharyngeal Edema</li> <li>Pruritus</li> <li>Superinfection</li> <li>Urticaria</li> <li>Pigment Dispersion Syndrome</li> </ul>	<ul style="list-style-type: none"> <li>Burning Sensation of Eyes</li> <li>Stinging of Eyes</li> <li>Conjunctival Hyperemia</li> <li>Corneal Edema</li> <li>Corneal Erosion</li> <li>Corneal Infiltrates</li> <li>Corneal Perforation</li> <li>Corneal Thinning</li> <li>Eye Pain and Irritation</li> <li>Headache</li> <li>Hypercreativity Reaction</li> <li>Iritis</li> <li>Ocular Edema</li> <li>Ophthalmic Inflammation</li> <li>Superficial Eye Infection</li> <li>Superficial Keratitis</li> <li>Tissue Damage</li> </ul>	
<b>Potential Contraindications/Precautions</b>	<ul style="list-style-type: none"> <li>Infants and patients less than 12 Years of age</li> <li>Lactation</li> <li>Pregnancy</li> </ul>	<ul style="list-style-type: none"> <li>Infants</li> <li>Lactation</li> <li>Pregnancy</li> </ul>	<ul style="list-style-type: none"> <li>Infants and patients less than 12 Years of age</li> <li>Lactation</li> <li>Pregnancy</li> </ul>	
<b>Storage</b>	Store at Room Temperature	20°C to 25°C (68°F to 77°F)		
<b>Auxiliary Labeling</b>	Protect from light	Store upright in original bottle	For professional use only	