

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

ImprimisRx is a registered trademark of Harrow Health, Inc.
©2020 Imprimis Rx. All Rights Reserved.
IMPO0312Rev3

**Questions?
Contact us at:
(844) 446-6979**

imprimis Rx®

Product Name:

Tropicamide
Proparacaine HCl
Phenylephrine HCl
Ketorolac Tromethamine

Ophthalmic Solution for Topical Use

ImprimisRx®
12264 El Camino Real, Suite 350
San Diego, CA 92130

	Tropicamide	Proparacaine HCl	Phenylephrine HCl	Ketorolac Tromethamine
Possible Indications	<ul style="list-style-type: none"> Cycloplegia Mydriasis 	<ul style="list-style-type: none"> Ocular Anesthesia 	<ul style="list-style-type: none"> Mydriasis Posterior Synechiae Uveitis with Posterior Synechiae 	<ul style="list-style-type: none"> Cystoid Macular Edema Miosis Postoperative Eye Pain Postoperative Ophthalmic Inflammation Seasonal Allergic Conjunctivitis Transient Burning or Stinging in the Eyes
Possible Adverse Effects	<ul style="list-style-type: none"> Blurred Vision Cardiorespiratory Arrest Headache Hypersensitivity Reaction Increased Intraocular Pressure Muscle Rigidity Nausea Pallor Photophobia Psychotic Reaction Superficial Punctate Keratitis Tachycardia Transient Burning or Stinging in the Eyes Vomiting Xerostomia 	<ul style="list-style-type: none"> Blurred Vision Stinging of the Eyes Hypersensitivity Reaction 	<ul style="list-style-type: none"> Acute Keratitis Acute Myocardial Infarction Allergic Conjunctivitis Blurred Vision Dermatitis Diaphoresis Dizziness Eye Irritation Headache Hypertension Lacrimation Pallor Photophobia Pigment Granules Liberation from the Iris Subarachnoid Hemorrhage Syncope Tachycardia Tremor Ventricular Premature contractions 	<ul style="list-style-type: none"> Burning Sensation of Eyes Stinging of Eyes Eye Irritation Hypersensitivity Pruritus and Skin Rash Superficial Keratitis Superficial Bacterial Eye Infection Conjunctival Hyperemia Corneal Edema Corneal Erosion Corneal Infiltrates Corneal Perforation Corneal Thinning Eye Pain Headache Iritis Ocular Edema Superficial Keratitis Tissue Damage
Potential Contraindications/Precautions	<ul style="list-style-type: none"> Patients less than 12 Years of age Lactation Pregnancy 	<ul style="list-style-type: none"> Infants Lactation Pregnancy 	<ul style="list-style-type: none"> Neonates and patients less than 12 years of age Lactation Pregnancy 	<ul style="list-style-type: none"> Patients less than 3 years of age Lactation Pregnancy Child bearing aged females
Storage	Store at Room Temperature	20°C to 25°C (68°F to 77°F)		
Auxiliary Labeling	Store upright in original bottle	Protect from light	For professional use only	