

The enclosed prescription medication is made by ImprimisRx® a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication.

This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

Questions?
Contact us at:
(844) 446-6979

imprimis Rx®

Product Name:

Prednisolone Acetate

Gatifloxacin

Bromfenac

Ophthalmic Suspension for Topical Use

	Prednisolone Acetate	Gatifloxacin	Bromfenac	
Possible Indications	Postoperative Ophthalmic Inflammation Postoperative Eye Pain	Bacterial Conjunctivitis	Postoperative Ophthalmic Inflammation Postoperative Ophthalmic Pain	
Possible Adverse Effects	Accommodation Disturbance Anterior Uveitis Blepharitis Conjunctival Hyperemia Conjunctivitis Corneal Perforation or Ulcer Decreased Visual Acuity Eye Infection Glaucoma Increased Intraocular Pressure Keratitis Mydriasis Optic Nerve Damage Visual Field Defect Wound Healing Impairment	Conjunctival Irritation Decreased Visual Acuity Dysgeusia Eye Discharge Eye Irritation Eye Pain Eye Redness Eyelid Edema Headache Increased Lacrimation Keratitis Papillary Conjunctivitis	Abnormal Sensation in Eyes Anaphylaxis Anterior Chamber Eye Hemorrhage Burning Sensation Conjunctival Hyperemia Corneal Erosion Corneal Perforation Corneal Thinning Corneal Ulcer Epithelial Keratopathy Eye Irritation Eye Pain Eye Pruritus Eye Redness Headache Hypersensitivity Reaction Iritis Keratitis Prolonged Bleeding Stinging Sensation	
Potential Contraindications/ Precautions	Infants Lactation Pregnancy Cataract Eye Infection Increased Intraocular Pressure	Infants Lactation Pregnancy	Lactation Pediatric Patients Pregnancy Corneal Disease Thrombocytopenia Rheumatoid Arthritis	
Storage	Store at Room Temperature	20°C to 25°C (68°F to 77°F)		
Auxiliary Labeling	Protect from light	Store upright in original bottle		