



**Product Name:**

Prednisolone Acetate

Moxifloxacin

Nepafenac

---

Ophthalmic Suspension for Topical Use

**Questions?  
Contact us at:  
(844) 446-6979**

ImprimisRx®  
12264 El Camino Real, Suite 350  
San Diego, CA 92130

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

		<b>Prednisolone Acetate</b>	<b>Moxifloxacin</b>	<b>Nepafenac</b>
	<b>Possible Indications</b>	Postoperative Ophthalmic Inflammation Postoperative Eye Pain	Bacterial Conjunctivitis	Postoperative Ophthalmic Inflammation Postoperative Ophthalmic Pain
	<b>Possible Adverse Effects</b>	Accommodation Disturbance Anterior Uveitis Blepharitis Conjunctival Hyperemia Conjunctivitis Corneal Perforation or Ulcer Decreased Visual Acuity Eye Infection Glaucoma Increased Intraocular Pressure Keratitis Mydriasis Optic Nerve Damage Visual Field Defect Wound Healing Impairment	Conjunctival Irritation Decreased Visual Acuity Dysgeusia Eye Discharge Eye Irritation Eye Pain Eye Redness Eyelid Edema Headache Increased Lacrimation Keratitis Papillary Conjunctivitis Pyrexia Subconjunctival Hemorrhage Tearing Hypersensitivity Reaction	Abnormal Sensation in Eyes Increased Intraocular Pressure Anterior Chamber Eye Hemorrhage Burning Sensation Conjunctival Hyperemia Corneal Erosion Corneal Perforation Corneal Thinning Corneal Ulcer Epithelial Keratopathy Eye Irritation Eye Pain Eye Pruritus Eye Redness Headache Hypersensitivity Reaction Iritis Keratitis Prolonged Bleeding Stinging Sensation
	<b>Potential Contraindications/ Precautions</b>	Infants Lactation Pregnancy Cataract Eye Infection Increased Intraocular Pressure	Infants Lactation Pregnancy	Lactation Pediatric Patients Pregnancy Corneal Disease Thrombocytopenia (bleeding disorders)
	<b>Storage</b>	Store at Room Temperature	20°C to 25°C (68°F to 77°F)	
	<b>Auxiliary Labeling</b>	Shake Well Before Use!	Store upright in original bottle	Protect from light.