

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs.

This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions.

ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

ImprimisRx is a registered trademark of Harrow Health, Inc. ©2020 ImprimisRx. All rights reserved. IMPO0376 Rev2

**Questions?
Contact us at:
(844) 446-6979**

ImprimisRx®
12264 El Camino Real, Suite 350
San Diego, CA 92130

imprimis Rx®

Product Name:

Prednisolone Acetate
Nepafenac

Ophthalmic Suspension for Topical Use

Prednisolone Acetate**Nepafenac****Possible Indications**

Postoperative Ophthalmic Inflammation
Postoperative Eye Pain

Postoperative Ophthalmic Inflammation
Postoperative Ophthalmic Pain

Possible Adverse Effects

Accommodation Disturbance
Anterior Uveitis
Blepharitis
Conjunctival Hyperemia
Conjunctivitis
Corneal Perforation or Ulcer
Decreased Visual Acuity
Eye Infection
Glaucoma
Increased Intraocular Pressure
Keratitis
Mydriasis
Optic Nerve Damage
Visual Field Defect
Wound Healing Impairment

Abnormal Sensation in Eyes
Increased Intraocular Pressure
Anterior Chamber Eye Hemorrhage
Burning Sensation
Conjunctival Hyperemia
Corneal Erosion
Corneal Perforation
Corneal Thinning
Corneal Ulcer
Epithelial Keratopathy
Eye Irritation
Eye Pain
Eye Pruritus
Eye Redness
Headache
Hypersensitivity Reaction
Iritis
Keratitis
Prolonged Bleeding
Stinging Sensation

**Potential Contraindications/
Precautions**

Infants
Lactation
Pregnancy
Cataract
Eye Infection
Increased Intraocular Pressure

Lactation
Pediatric Patients
Pregnancy
Corneal Disease
Thrombocytopenia (bleeding disorders)

Storage

Store at Room Temperature

20°C to 25°C (68°F to 77°F)

Auxiliary Labeling

Shake Well Before Use!