

The enclosed prescription medication is made by ImprimisRx, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs.

This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions.

ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

**Questions?  
Contact us at:  
(844) 446-6979**

**imprimis** Rx<sup>®</sup>

**Product Name:**

**Timolol  
Latanoprost**

---

Ophthalmic Solution  
Preservative-Free

	<b>Timolol Maleate</b>	<b>Latanoprost</b>	
<b>Possible Indications</b>	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma	
<b>Possible Adverse Effects</b>	<p>Ocular burning/stinging on instillation</p> <p>Dry eyes</p> <p>Itching</p> <p>Foreign body sensation</p> <p>Discomfort in the eye</p> <p>Eyelid erythema</p> <p>Conjunctival injection</p> <p>Headache</p> <p>Eye pain</p> <p>Epiphora</p> <p>Photophobia</p> <p>Blurred or abnormal vision</p> <p>Corneal fluorescein staining</p> <p>Keratitis</p> <p>Blepharitis</p> <p>Cataract</p> <p>Cardiac arrhythmia, palpitation, bradycardia, Syncope, heart block, cerebral vascular accident, cerebral ischemia, cardiac failure and cardiac arrest</p>	<p>Iris pigmentation changes</p> <p>Eyelid skin darkening</p> <p>Eyelash changes (increased length, thickness, pigmentation and number of lashes)</p> <p>Iritis/uveitis</p> <p>Macular edema, including cystoid macular edema</p> <p>Foreign body sensation</p> <p>Punctate keratitis</p> <p>Stinging</p> <p>Conjunctival hyperemia</p> <p>Blurred vision</p> <p>Itching</p> <p>Burning</p> <p>Increased pigmentation of the iris</p> <p>Excessive tearing</p> <p>Eyelid discomfort/pain</p> <p>Dry eye</p> <p>Eye pain</p> <p>Eyelid margin crusting</p> <p>Erythema of the eyelid</p> <p>Photophobia</p>	
<b>Potential Contraindications/Precautions</b>	<p>Patients Less than 12 Years</p> <p>Lactation</p> <p>Pregnancy</p> <p>Bronchial Asthma or History Of.</p> <p>Sinus Bradycardia, Second or Third Degree Atrioventricular Block; Overt Heart Failure; Cardiogenic Shock.</p>	<p>Patients Less than 3 Years</p> <p>Lactation</p> <p>Pregnancy</p> <p>Child Bearing Aged Females</p>	
<b>Storage</b>	Store under refrigeration	2°C to 8°C (36°F to 46°F)	Protect from light.
<b>Auxiliary Labeling</b>	For professional use only.	Store upright in original bottle.	