

The enclosed prescription medication is made by ImprimisRx, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs.

This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions.

ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

Questions?
Contact us at:
(844) 446-6979

imprimis Rx[®]

Product Name:

**Timolol
Dorzolamide
Latanoprost**

Ophthalmic Solution
Preservative-Free

	Timolol Maleate	Dorzolamide	Latanoprost
Possible Indications	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma
Possible Adverse Effects	Ocular burning/stinging on instillation Dry eyes Itching Foreign body sensation Discomfort in the eye Eyelid erythema Conjunctival injection Headache Eye pain Epiphora Photophobia Blurred or abnormal vision Corneal fluorescein staining Keratitis Blepharitis Cataract Cardiac arrhythmia, palpitation, bradycardia, Syncope, heart block, cerebral vascular accident, cerebral ischemia, cardiac failure and cardiac arrest	Ocular burning, stinging, or discomfort immediately following ocular administration Bitter taste Superficial punctate keratitis Ocular allergic reaction Conjunctivitis and lid reactions Blurred vision Eye redness Tearing Dryness Photophobia Headache Nausea Asthenia/fatigue Skin rashes Urolithiasis Iridocyclitis	Iris pigmentation changes Eyelid skin darkening Eyelash changes (increased length, thickness, pigmentation and number of lashes) Iritis/uveitis Macular edema, including cystoid macular edema Foreign body sensation Punctate keratitis Stinging Conjunctival hyperemia Blurred vision Itching Burning Increased pigmentation of the iris Excessive tearing Eyelid discomfort/pain Dry eye Eye pain Eyelid margin crusting Erythema of the eyelid Photophobia
Potential Contraindications/Precautions	Patients Less than 12 Years Lactation Pregnancy Bronchial Asthma or History Of. Sinus Bradycardia, Second or Third Degree Atrioventricular Block; Overt Heart Failure; Cardiogenic Shock.	Neonates and Patients Less than 12 Years Lactation Pregnancy	Patients Less than 3 Years Lactation Pregnancy Child Bearing Aged Females
Storage	Store under refrigeration	2°C to 8°C (36°F to 46°F)	Protect from light.
Auxiliary Labeling	For professional use only.	Store upright in original bottle.	