

The enclosed prescription medication is made by ImprimisRx, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs.

This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions.

ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

ImprimisRx is trademarks of Harrow Health, Inc. ©2019 ImprimisRx. All rights reserved.  
IMPO0410 Rev2 11/19

**Questions?  
Contact us at:  
(844) 446-6979**

ImprimisRx  
12264 El Camino Real, Suite 350  
San Diego, CA 92130

**imprimis** Rx<sup>®</sup>

**Product Name:**

**Timolol  
Brimonidine  
Dorzolamide**

---

Ophthalmic Solution  
Preservative-Free

		<b>Timolol Maleate</b>	<b>Brimonidine Tartrate</b>	<b>Dorzolamide</b>
	<b>Possible Indications</b>	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma	Treatment of elevated intraocular pressure in patients with ocular hypertension or open-angle glaucoma
	<b>Possible Adverse Effects</b>	Ocular burning/stinging on instillation Dry eyes Itching Foreign body sensation Discomfort in the eye Eyelid erythema Conjunctival injection Headache Eye pain Epiphora Photophobia Blurred or abnormal vision Corneal fluorescein staining Keratitis Blepharitis Cataract Cardiac arrhythmia, palpitation, bradycardia, Syncope, heart block, cerebral vascular accident, cerebral ischemia, cardiac failure and cardiac arrest	Oral dryness Ocular hyperemia Burning and Stinging on Instillation Headache Blurred vision Foreign body sensation Fatigue/drowsiness Conjunctival follicles Ocular allergic reactions Ocular pruritis Corneal staining/erosion Photophobia Eyelid erythema Ocular ache/pain Upper respiratory symptoms Dizziness Blepharitis, ocular irritation Lid crusting Conjunctival hemorrhaging Abnormal taste, insomnia, depression, hypertension, anxiety, palpitations/arrhythmias, nasal dryness and syncope	Ocular burning, stinging, or discomfort immediately following ocular administration Bitter taste Superficial punctate keratitis Ocular allergic reaction Conjunctivitis and lid reactions Blurred vision Eye redness Tearing Dryness Photophobia Headache Nausea Asthenia/fatigue Skin rashes Urolithiasis Iridocyclitis
	<b>Potential Contraindications/Precautions</b>	Patients Less than 12 Years Lactation Pregnancy Bronchial Asthma or History Of. Sinus Bradycardia, Second or Third Degree Atrioventricular Block; Overt Heart Failure; Cardiogenic Shock.	Infants Lactation Pregnancy Concomitant MAO Inhibitor Therapy	Neonates and Patients Less than 12 Years Lactation Pregnancy
	<b>Storage</b>	Store at room temperature	20°C to 25°C (68°F to 77°F)	
	<b>Auxiliary Labeling</b>	For professional use only.	Store upright in original bottle.	Protect from light.