



Product Name:

Klarity Drops
(Glycerin and dextran-based vehicle)

Ophthalmic Solution for Topical Use
Preservative-Free

**Questions?
Contact us at:
(844) 446-6979**

ImprimisRx®
12264 El Camino Real, Suite 350
San Diego, CA 92130

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

Glycerin

Dextran

Possible Indications

Ophthalmic lubricant; for temporary relief of burning and eye irritation due to dry eyes

Protect and lubricate mucous membrane surfaces and relieve dryness and irritation

Possible Adverse Effects

Blurred vision
Crusting of eyelid
Stinging of eyes (mild)

Mild eye burning or irritation
Itching or redness of your eye
Watery eyes
Blurred vision

Potential Contraindications/ Precautions

Self-medication:
When used for self-medication, stop use and contact healthcare provider if changes in vision, eye pain, continued redness or irritation occur, or if the condition worsens or persists for more than 72 hours. To avoid contamination, do not touch tip of container to any surface. Replace cap after using. Do not use if solution is cloudy or changes color.

Self-medication:
When used for self-medication, stop use and contact healthcare provider if changes in vision, eye pain, continued redness or irritation occur, or if the condition worsens or persists for more than 72 hours. To avoid contamination, do not touch tip of container to any surface. Replace cap after using. Do not use if solution is cloudy or changes color.

Storage

Store at Room Temperature

20°C to 25°C (68°F to 77°F)

Auxiliary Labeling

Store upright in original bottle

FOR PRESCRIPTION USE ONLY