



**Product Name:**

Tobramycin – Vancomycin

---

Fortified Ophthalmic Solution

**Questions?  
Contact us at:  
(844) 446-6979**

ImprimisRx®  
1000 Aviara Parkway, Suite 220  
Carlsbad, CA 92011

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound essentially copies of commercially available products. References available upon request.

## Tobramycin

## Vancomycin

### Possible Indications

Treatment of infections of the eye caused by susceptible bacteria

Treatment of infections of the eye caused by susceptible bacteria

### Possible Adverse Effects

Ocular burning/stinging on instillation  
Lid itching and swelling  
Conjunctival erythema  
Anaphylactic reaction  
Stevens-Johnson syndrome  
Erythema multiforme

Ocular burning/stinging on instillation  
Lid itching and swelling  
Conjunctival erythema  
Anaphylactic reaction  
Stevens-Johnson syndrome

The following are adverse reactions reported with systemic aminoglycosides:

- Neurotoxicity
- Ototoxicity
- Nephrotoxicity
- Aminoglycosides may aggravate muscle weakness in patients with known or suspected neuromuscular disorders, such as myasthenia gravis or Parkinson's disease.

The following are adverse reactions reported with systemic vancomycin administration:

- Nephrotoxicity
- Onset of pseudomembranous colitis symptoms
- Ototoxicity
- Neutropenia

Hemorrhagic occlusive retinal vasculitis (HORV) may be associated with the use of intraocular vancomycin (including intravitreal, intracameral bolus, and irrigating solution containing vancomycin). Although the precise etiology of HORV remains unproven, each surgeon should weigh the potential risk of vancomycin against the risk of endophthalmitis.

### Potential Contraindications/ Precautions

Patients Less than 12 Years  
Nursing Mothers  
Pregnancy

Patients Less than 12 Years  
Nursing Mothers  
Pregnancy

### Storage

Store under refrigeration  
2°C to 8°C (36°F to 46°F)

Store under refrigeration  
2°C to 8°C (36°F to 46°F)

### Auxiliary Labeling

Protect from light

Protect from light