

The enclosed prescription medication is made by ImprimisRx® a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. This document is for professional use only. This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication.

This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound copies of commercially available products. References available upon request.

ImprimisRx is a registered trademark of Imprimis Pharmaceuticals, Inc.
©2018 Imprimis Pharmaceuticals, Inc. All Rights Reserved. IMPO0318 9/18

Questions?
Contact us at:
(844) 446-6979

Imprimis Pharmaceuticals, Inc.
12264 El Camino Real, Suite 350
San Diego, CA 92130

Imprimis Rx®

Product Name:

Prednisolone Acetate

Gatifloxacin

Bromfenac

Ophthalmic Suspension for Topical Use

	Prednisolone Acetate	Gatifloxacin	Bromfenac	
Possible Indications	<ul style="list-style-type: none"> Postoperative Ophthalmic Inflammation Postoperative Eye Pain 	<ul style="list-style-type: none"> Bacterial Conjunctivitis 	<ul style="list-style-type: none"> Postoperative Ophthalmic Inflammation Postoperative Ophthalmic Pain 	
Possible Adverse Effects	<ul style="list-style-type: none"> Burning or Stinging Sensation Accommodation Disturbance Conjunctival Hyperemia Decreased Visual Acuity Eye Infection Glaucoma Increased Intraocular Pressure Keratitis Mydriasis Optic Nerve Damage Subscapular Posterior Cataract Visual Field Defect Wound Healing Impairment Corneal Ulcer Viral Infection 	<ul style="list-style-type: none"> Conjunctival Irritation Decreased Visual Acuity Dysgeusia Eye Discharge Eye Irritation Eye Pain Eye Redness Eyelid Edema Headache Increased Lacrimation Keratitis Papillary Conjunctivitis 	<ul style="list-style-type: none"> Abnormal Sensation in Eyes Anaphylaxis Anterior Chamber Eye Hemorrhage Burning Sensation Conjunctival Hyperemia Corneal Erosion Corneal Perforation Corneal Thinning Corneal Ulcer Epithelial Keratopathy Eye Irritation Eye Pain Eye Pruritus Eye Redness Headache Hypersensitivity Reaction Iritis Keratitis Prolonged Bleeding Stinging Sensation 	
Potential Contraindications/Precautions	<ul style="list-style-type: none"> Infants Lactation Pregnancy Cataract Eye Infection Increased Intraocular Pressure 	<ul style="list-style-type: none"> Infants Lactation Pregnancy 	<ul style="list-style-type: none"> Lactation Pediatric Patients Pregnancy Corneal Disease Thrombocytopenia Rheumatoid Arthritis 	
Storage	<ul style="list-style-type: none"> Store at Room Temperature 	<ul style="list-style-type: none"> 20°C to 25°C (68°F to 77°F) 		
Auxiliary Labeling	<ul style="list-style-type: none"> Protect from light 	<ul style="list-style-type: none"> Store upright in original bottle 		