

Formulation Description: Triamcinolone Acetonide and Moxifloxacin HCl Intraocular Suspension for Injection

The enclosed prescription medication is made by ImprimisRx®, a pharmaceutical compounding company that specializes in customizing medications to meet unique patient and practitioner needs. **This document is for professional use only.** This document contains information related to the individual ingredients contained in the above described compounded formulation. Use of this formulation has not been reviewed by the U.S. Food and Drug Administration (FDA) for any indication. This information may not cover all possible uses, directions, side effects, precautions, allergic reactions or drug interactions. ImprimisRx dispenses compounded medicines to licensed practitioners and individually identified patients with valid prescriptions. Compounded medications are not reviewed by the FDA for safety or efficacy. ImprimisRx does not compound copies of commercially available products. References available upon request.

	Triamcinolone Acetonide	Moxifloxacin HCl
Possible Indications	Postoperative Ophthalmic Inflammation Sympathetic Ophthalmia Diffuse Posterior Uveitis Allergic Corneal Margin Ulcer Iritis and Keratitis Optic Neuritis	Bacterial Conjunctivitis
Possible Adverse Effects	Anaphylactoid Shock Blindness Cataract Corneal Perforation Cortical Cataract Conjunctival Hemorrhage Eye Pain, Infection, Irritation Pruritus Abnormal Sensation in Eyes Abnormal Sensory Symptoms Edema Bleeding at Injection Site Glaucoma Hypertension Hypopyon Increased Intraocular Pressure Increased Lacrimation Vitreous Detachment Vitreous Opacity Wound Healing Impairment	Conjunctival Irritation Fever Eye Irritation Airway Obstruction Anaphylaxis Angioedema Circulatory Shock Dyspnea Facial Edema Fungal Superinfection Laryngeal Edema Loss of Consciousness Pharyngeal Edema Pruritus Superinfection Urticaria Pigment Dispersion Syndrome
Potential Contraindications/Precautions	Infants Lactation Pregnancy Cataract Eye Infection Increased Intraocular Pressure	Infants Lactation Pregnancy
Storage	Store at room temperature.	20°C to 25°C
Auxiliary Labeling	Shake well before use! For professional use only.	Store upright in original bottle.